

The functional role of sodium taurocholate co-transporting polypeptide NTCP in the life cycle of hepatitis B, C and D viruses

Carla Eller^{1,2}, Laura Heydmann^{1,2}, Che C. Colpitts³, Eloi R. Verrier^{1,2}, Catherine Schuster^{1,2},
Thomas F. Baumert^{1,2,4*}

¹Inserm, U1110, Institut de Recherche sur les Maladies Virales et Hépatiques, 67000 Strasbourg, France;

²Université de Strasbourg, 67000 Strasbourg, France; ³Division of Infection and Immunity, University College London, London, United Kingdom; ⁴Institut Hospitalo-Universitaire, Pôle Hépatodigestif, Nouvel Hôpital Civil, 67000 Strasbourg, France.

***Corresponding author:** Prof. Thomas F. Baumert, MD; Inserm U1110, Institut de Recherche sur les Maladies Virales et Hépatiques, Université de Strasbourg, 3 Rue Koeberlé, 67000 Strasbourg, France; Phone: +33 3 68 85 37 03, Fax: +33 3 68 85 37 24, e-mail: thomas.baumert@unistra.fr

ORCID of the authors: 0000-0001-7835-9468, 0000-0002-2273-2267, 0000-0003-2474-1834, 0000-0002-9022-5611, 0000-0001-7281-4511, 0000-0002-8864-2168

Acknowledgments / Funding

This work was supported by Inserm, the University of Strasbourg, the European Union (ERC-2014-AdG-671231-HEPCIR, Infect-ERA hepBccc, EU H2020 Hep-CAR 667273), ANRS (2015/1099), the French Cancer Agency (ARC IHU201301187) and the National Institute of Allergy and Infectious Diseases of the National Institutes of Health under award number R03AI131066. CCC acknowledges fellowships from the Canadian Institutes of Health Research (201411MFE-338606-245517) and the Canadian Network on Hepatitis C. ERV is the recipient of an ANRS fellowship (ECTZ50121).

Abstract

Chronic hepatitis B, C and D virus (HBV, HCV and HDV) infections are a major cause of liver disease and cancer worldwide. Despite employing distinct replication strategies, the three viruses are exclusively hepatotropic and therefore depend on hepatocyte-specific host factors. The sodium taurocholate co-transporting polypeptide (NTCP), a transmembrane protein highly expressed in human hepatocytes that mediates the transport of bile acids, plays a key role in HBV and HDV entry into hepatocytes. Recently, NTCP has been shown to modulate HCV infection of hepatocytes by regulating innate antiviral immune responses in the liver. Here we review the current knowledge of the functional role and the molecular and cellular biology of NTCP in the life cycle of the three major hepatotropic viruses, highlight the impact of NTCP as an antiviral target and discuss future avenues of research.

Keywords: Liver cell biology, bile acid transport, host factor, anti-viral therapy, hepatocytes.

Introduction

Every year, viral hepatitis is estimated to cause around 1.3 million deaths worldwide, mainly through chronic liver disease and hepatocellular carcinoma (HCC). Approximately 95% of these deaths are caused by hepatitis B and C viruses (HBV, HCV) [1]. Despite the availability of an effective vaccine for HBV, 250 million people are chronically infected by the virus worldwide [2]. An estimated 5% of HBV patients are co-infected with hepatitis D virus (HDV), a satellite virus hijacking HBV envelope proteins to assemble its infectious viral particles. HDV co-infection worsens the outcome of HBV infection and treatment of HBV-HDV co-infected patients is less effective [3, 4]. Moreover, around 70 million people are living with chronic HCV infection and, despite the existence of effective curative strategies, the incidence of HCV is still increasing [3].

Remarkable progress has recently been made for treatment of HCV infection. The development and approval of direct acting antivirals (DAAs) specifically targeting viral proteins now allows for HCV cure, but these therapies remain inaccessible for the majority of HCV patients [5]. For chronic HBV infection, two therapeutic approaches are used to suppress viral replication: pegylated interferon and nucleos(t)ide analogues (NUCs). While these treatments allow control of HBV infection, viral eradication is rare and, in most cases, lifelong therapy is required [6]. For patients with chronic HBV/HDV co-infection, the current treatment options are limited to interferon-alpha (IFN α) and its pegylated derivative. Furthermore, although current antivirals decrease the risk of HCC, they are not sufficient to eliminate the risk [7, 8]. In order to effectively combat these hepatotropic viruses,

it is necessary to improve existing therapies and uncover new strategies for prevention and treatment of viral hepatitis.

Alternative strategies against chronic HBV and HCV infection include host-targeting agents (HTA), which target host factors required for viral replication. HTAs have been shown to be promising candidates for the prevention and treatment of infections by various pathogens, including HBV and HCV [9–11]. This approach requires a profound understanding of the viral life cycle and the virus-host interactions involved. Indeed, the identification of the human sodium taurocholate co-transporting polypeptide (NTCP) as a functional receptor for HBV/HDV infection [12, 13] opened perspectives for new antiviral strategies. Several entry inhibitors for treatment of HBV infection targeting NTCP are now in development [14–19]. Furthermore, this crucial discovery has allowed the development of novel infectious model systems that will enable an improved understanding of the complete HBV/HDV viral life cycle [20]. However, the regulatory role of NTCP in HCV host cell infection, and its potential immunomodulatory activities in hepatocytes, should not be overlooked. The aim of this review is to summarize what is known about the interactions of NTCP with three major hepatitis viruses during infection, to describe the molecular mechanisms, and to highlight possible applications in research and therapy.

Sodium taurocholate co-transporting polypeptide, a bile acid transporter

The circulation of bile and bile components between human intestine enterocytes and liver parenchymal cells is known as the enterohepatic circulation (EHC) [21]. In the liver, bile acids are mainly involved in cholesterol metabolism and elimination of toxic compounds [22]. Interestingly, bile acids have also been shown to inhibit interferon (IFN) signaling pathways, resulting in reduced expression of IFN-stimulated genes (ISG) [23, 24]. In hepatocytes, bile acid homeostasis is maintained by the interplay between uptake, synthesis and secretion of bile acids. The major hepatic uptake transporter for conjugated bile acids in humans is sodium taurocholate co-transporting polypeptide (NTCP) [25]. NTCP is predominantly expressed at the hepatic basolateral membrane and is involved in the recycling of bile acids from portal blood to hepatocytes in a sodium-dependent manner [21]. NTCP is a member of the solute carrier family SLC10 and is encoded by *SLC10A1* [26, 27]. *SLC10A1* mRNA is translated into a 349 amino acid glycosylated phosphoprotein with seven or nine transmembrane domains [21, 28–31]. While the exact function of some SLC10 family members remains unknown, all of them are thought to be sodium-dependent transporters [21]. Interestingly, bile acid transport through NTCP can be blocked by small molecules already in clinical use, such as cyclosporine A (CsA, an immunosuppressive drug used in transplantation) or ezetimibe (used for hypercholesterolemia) [16, 32].

Hepatic bile acid metabolism is tightly regulated, including at the transcriptional level (see Figure 1) [33]. Upon bile acid activation, the nuclear factor Farnesoid X Receptor (FXR) indirectly downregulates several target genes through transcriptional induction of the small heterodimer partner (SHP) [34, 35], including the first and rate-limiting enzyme in bile acid biosynthesis cholesterol 7 α -hydroxylase (CYP7A1) [36, 37]. FXR also directly activates the expression of the bile salt export pump (BSEP, ABCB11), which is expressed at the apical membrane and secretes conjugated bile acids into the bile canaliculus in an ATP-dependent manner [38, 39]. FXR does not directly interact with the promoter of human *SLC10A1* but induces the expression of different factors to indirectly repress *slc10a1* expression in rat and mouse, although mechanisms of transcriptional regulation of human NTCP remain unknown [40–42]. In hepatic inflammation, the cytokines tumor necrosis factor alpha (TNF- α), interleukin (IL)-1 β , and IL-6 downregulate mRNA levels of *SLC10A1* and reduce the transporter protein expression [43–45]. The downregulation of NTCP expression in the human liver has been implicated in several cholestasis pathologies. The reduction of NTCP expression could explain impaired hepatic bile acid uptake, resulting in cholestasis and jaundice. Several studies have shown a downregulation of bile salt transporters in primary biliary cirrhosis [46, 47]. Interestingly, a recent study showed a suppression of NTCP expression via cyclin D1 in hepatocellular carcinoma (HCC) [48]. These data may explain the low expression level of NTCP in HCC-derived cell lines, such as Huh7 and clones or HepG2.

The localization and membrane expression of NTCP is controlled by post-translational mechanisms [49]. For example, cyclic adenosine monophosphate (cAMP) plays a role in stimulating the dephosphorylation and membrane translocation of NTCP (see Figure 1) [50–52]. Sequencing analysis of NTCP revealed the existence of several ethnic-dependent single nucleotide polymorphisms (SNPs) which may alter NTCP activities [53]. For example, mutation S267F, found in 7.5% of allele frequencies in Chinese Americans, is associated with an almost complete loss of bile acid uptake function. However, no pathologies have been described resulting from these NTCP polymorphisms and their clinical roles remain controversial [54]. Besides its major role in the bile acid uptake system, Yan *et al.* described the crucial role of NTCP on HBV and HDV entry [12]. For the time being, NTCP remains the only described HBV and HDV entry receptor.

NTCP is a host factor for HBV/HDV infection

Hepatitis B virus is the prototypic member of the *Hepadnaviridae* family of small enveloped hepatotropic DNA viruses. Its envelope consists of three different forms of the HBV surface protein (HBsAg) – the small (S), middle (M) and large (L) proteins. Importantly, the preS1-domain of L envelope protein is known to bind the hepatocyte cell surface and is required for HBV and HDV entry [55]. The HBV capsid is comprised of HBV core protein (HBcAg) and encapsidates a partially double-stranded relaxed circular DNA (rcDNA) genome of 3.2 kilobases. Upon infection of hepatocytes, genomic rcDNA is converted into covalently closed circular DNA (cccDNA), a minichromosome-like structure that persists in the nucleus as a central transcription template for all viral RNAs [56]. The presence of cccDNA in the nucleus is thought to be responsible for viral rebound after withdrawal of NUC therapy that targets reverse transcription, a late step in the HBV life cycle. Therefore, removal of cccDNA from HBV-infected hepatocytes will be essential to achieve the goal of HBV cure [57].

HDV is a defective hepatotropic virus which depends on HBV surface proteins for assembly of infectious virions and viral entry [58]. The HDV genome is a negative single-stranded circular RNA of nearly 1700 nucleotides containing one functional open reading frame, which encodes the hepatitis delta protein (HDAg) expressed in small and large form. Replication of HDV RNA and transcription of HDAg mRNA in the nucleus depends on host cell polymerases, including DNA-dependent RNA polymerase II. Both forms of the delta protein are then produced and reimported in the nucleus where they bind to genomic RNA to form the ribonucleoprotein (RNP), which is then exported into the cytoplasm and is associated with HBV envelope proteins to form a mature HDV virion [59]. Thus, HDV enters hepatocytes using the same pathways as HBV, and depends on the same host factors for host cell binding and entry. HDV is therefore a useful surrogate model for HBV entry.

The first step of viral infection is virion binding to attachment factors and receptors at the host cell surface. This specific interaction between viral surface proteins and host entry receptors often determines the tissue tropism and host range of the virus [60]. HBV and its satellite virus HDV share HBV envelope proteins and are known to exclusively infect human, chimpanzee and tree shrew (*Tupaia belangerii*) hepatocytes, suggesting the involvement of species- and liver-specific cell surface factors in the common entry process of these viruses [20]. Two elements of the HBV envelope proteins are necessary for interaction with these factors. One determinant of infectivity resides in the surface-exposed cysteine-rich antigenic loop (AGL), a polypeptide located in the S domain of all three envelope proteins [61, 62]. The second known infectivity determinant is a receptor binding site in the N-terminal pre-S1 domain of the L-HBsAg [55]. This domain is post-translationally modified by addition of myristic

acid [63], and this myristoylation is essential for virion infectivity [64, 65]. A synthetic myristoylated peptide comprising the N-terminal amino acids 2 to 78 of the pre-S1 domain prevents HBV infection [66].

As for many viruses [67, 68], HBV/HDV infection requires the initial attachment to the glycosaminoglycan (GAG) side chains of heparan sulfate proteoglycans (HSPGs) [69]. Both the antigenic loop of all HBV envelope proteins and the preS1-region of HBsAg-L are involved in this interaction [69, 70]. Indeed, glypican-5 (GPC5), a member of the glypican family of HSPGs, acts as an entry factor for HBV and HDV (see Figure 2) [71]. After this initial step of HBV/HDV attachment to HSPGs, the virions bind to a high-affinity receptor via the preS1-domain [72], allowing uptake into hepatocytes. Despite the discovery of several preS1-interacting proteins that did not affect HBV infectivity [73–78], the identity of the HBV/HDV entry receptor remained unclear until 2012, when Yan *et al.* identified NTCP as a functional receptor for HBV and HDV infection. Using a labeled preS1 peptide as a bait in *Tupaia* hepatocytes, a mass spectrometry purification of preS1-bound proteins, and validation in human hepatocytes, it was shown that NTCP specifically interacts with the HBV receptor-binding domain preS1, allowing viral entry [12]. Zhong *et al.* showed that *Tupaia* NTCP mediates entry of woolly monkey HBV, indicating that NTCP orthologs act as a common cellular receptor for known primate hepadnaviruses [79]. Differential gene expression patterns between non-susceptible undifferentiated and susceptible differentiated HepaRG cells validated the role of NTCP as a specific receptor for HBV and HDV [13]. Moreover, silencing of NTCP in primary *Tupaia* hepatocytes (PTH) or differentiated HepaRG cells inhibits HBV and HDV infection [12, 13]. Exogenous expression of NTCP directly renders non-susceptible hepatoma cell lines susceptible to HBV and HDV infection, while entry inhibitors derived from the preS1 peptide efficiently inhibit this infection [12]. In addition, the S267F mutant of NTCP, conferring a loss of bile acid uptake function, is significantly associated with resistance to chronic hepatitis B and decreased risk of cirrhosis and liver cancer development, supporting the role of NTCP as cellular receptor for HBV in human infection [80–82]. However, S267F homozygote patients can still be infected by HBV, suggesting the existence of alternative receptors allowing viral entry in the absence of functional NTCP [83].

Interestingly, expression of human (but not mouse) NTCP in non-susceptible hepatocarcinoma cells confers limited susceptibility to infection. For robust infection, addition of dimethyl sulfoxide (DMSO) to culture medium is essential [13]. The fact that human hepatoma cell lines HepG2 and Huh7 are not susceptible to HBV and HDV infection without exogenous expression of NTCP is consistent with reports that NTCP expression is reduced in human hepatocellular carcinoma cells [48, 84]. NTCP expression rapidly decreases over time following isolation of cultured PTHs, which supports observations that primary human hepatocytes (PHH) remain

susceptible to HBV infection *in vitro* only for a few days after isolation [12, 85]. Considering the predominant expression of NTCP in the liver, this receptor is likely to contribute to the hepatotropism of both viruses [12]. In addition, NTCP protein sequences vary among mammalian species, which might contribute to the narrow species tropism of HBV and HDV infection. For example, monkey NTCP does not support HBV and HDV infection despite a high protein sequence homology to human NTCP. Replacing amino acids 157–165 of nonfunctional monkey NTCP with the human counterpart conferred susceptibility to both HDV and HBV infection [12]. The fact that hepatocytes from cynomolgus and rhesus macaques and pigs become fully susceptible to HBV upon hNTCP expression indicates that NTCP is the key host factor limiting HBV infection in these species [86].

As a key host factor enabling HBV and HDV infection *in vitro*, the discovery of NTCP has been crucial for the development of novel animal models supporting virus infection. Indeed, only chimpanzees and *Tupaia* can experimentally support HBV and HDV infections [87]. The state-of-the-art mouse model for the study of HBV/HDV consists of liver-engrafted humanized chimeric uPa/SCID or FRG mice, which support virus entry and replication, but lack an efficient immune system, limiting the study of virus-host interactions [87]. The recent development of human NTCP-expressing transgenic mice opened perspectives for the development of novel immune-competent animal models for the investigation of HDV infection and HDV-induced pathogenesis *in vivo* [88]. As HBV infection is limited in mouse cells expressing hNTCP, probably due to the lack of a key host factor [89], it should be noted that hNTCP-transgenic mice are not susceptible to HBV infection. Recently, an elegant study demonstrated that vector-mediated expression of hNTCP in the hepatocytes of rhesus macaques conferred susceptibility to HBV infection, providing a robust and relevant model for the study of HBV infection, including its interaction with adaptive immunity and the understanding of viral clearance [90].

Overall, NTCP was identified as the long-sought preS1-specific HBV receptor contributing to HBV liver tropism and species specificity [13]. Targeting the interactions between the HBV preS1-domain and its receptor NTCP required for HBV/HDV entry is a promising strategy to block viral entry for both viruses.

NTCP as a therapeutic target for HBV/HDV infection

Even before the identification of NTCP as HBV/HDV receptor, entry inhibitors derived from the HBV preS1 were shown to efficiently inhibit HBV infection *in vitro* and *in vivo* [91, 92]. One of these compounds, the myristoylated preS1-derived peptide (also called Myrcludex B or MyrB), efficiently prevents HBV dissemination *in vivo* and hinders amplification of the cccDNA pool in infected human hepatocytes [14]. MyrB is the first HBV/HDV entry inhibitor targeting NTCP to reach clinical trials [93], where it was shown to have a good safety profile with a mild

and reversible elevation of serum bile acid salts [93, 94]. Phase IIa clinical studies revealed a marked antiviral effect of MyrB, as measured by HDV RNA, HBV DNA and improvement of biochemical disease activity (ALT), when used in combination with IFN therapy, although there was no significant decrease in HBsAg levels. In monotherapy, however, MyrB did not show significant antiviral activity [94]. Further studies are necessary to confirm these results obtained in small patient cohorts [95].

Importantly, the identification of NTCP as the first HBV/HDV entry receptor has accelerated the discovery and development of several new potential entry inhibitors. Binding of myristoylated preS1-derived peptide to NTCP was shown to interfere with the physiological bile acid transport function of NTCP, indicating that NTCP-inhibiting drugs might be able to block HBV infection [96]. In a study evaluating FDA approved therapeutics with documented inhibitory effect on NTCP cellular function against HDV entry, three of these molecules (irbesartan, ezetimibe, and ritonavir) inhibited HDV infection *in vitro* [97]. The inhibitory effect of ezetimibe on HBV infection had already been described previously without understanding its interactions with NTCP [98]. In 2014, Watashi *et al.* evaluated the effect of compounds on the early phase of the HBV life cycle to identify cyclosporine A as an HBV entry inhibitor targeting NTCP [15]. In the same year, Nkongolo *et al.* characterized the effect of cyclosporine A, a cholestasis-inducing drug inhibiting NTCP bile acid transport [32, 97, 98], against HBV/HDV infection and found that inhibition of entry resulted from interference with the NTCP receptor [16]. The screening of FDA/EMA-approved drugs or small molecules for interaction with NTCP allowed the identification of several additional potential HBV/HDV entry inhibitors targeting NTCP [18, 19]. All of these NTCP-targeting HBV/HDV entry inhibitors concomitantly inhibit the transporter function of NTCP and impair bile acid uptake into hepatocytes, increasing the risk of adverse effects. NTCP-deficient mice and a patient with NTCP deficiency were shown to exhibit an elevated level of serum bile acids and to develop related pathologies including growth retardation and hypercholanemia [101, 102].

Two different strategies to selectively inhibit HBV entry without impairing bile acid uptake have been suggested recently. Shimura *et al.* showed that cyclosporine A derivatives SCY450 and SCY995 inhibit HBV/HDV entry without interfering with the NTCP transporter activity (see Figure 2) [17]. Tsukuda *et al.* identified an oligomeric flavonoid, proanthocyanidin (PAC) and its analogs, as a new class of entry inhibitors, which directly target the preS1-domain of the HBV large envelope protein and thereby prevent its attachment to NTCP. By directly targeting HBV particles, PAC impairs HBV infectivity without affecting the NTCP-mediated bile acid transport activity [103]. Further studies are required to determine if these novel inhibitory strategies will show efficacy *in vivo* and in clinical studies in co-treatment with NUC therapy.

NTCP is a host factor for HCV infection

Hepatitis C virus is an enveloped single-stranded positive-sense RNA virus in the *Flaviviridae* family. The host-cell derived lipid envelope contains the two viral envelope glycoproteins, E1 and E2 [104]. Within the envelope, an icosahedral capsid contains the RNA genome of 9.6 kilobases. Like HBV and HDV, attachment of HCV to hepatocytes is mediated by HPSGs on the host cell surface [105–107]. Following attachment, the envelope glycoprotein E2 mediates interactions with a series of specific cellular entry factors, including CD81 and claudin-1 (see Figure 2) [108–111]. HCV is internalized via endocytosis in a clathrin- and dynamin-dependent process [112]. Following fusion with early endosomal membranes, the HCV genome is released into the cytosol, where it is translated into a polyprotein cleaved by viral and host proteases. The HCV genome is replicated directly into RNA without passing through a DNA intermediate [113]. Therefore, HCV entry and replication steps are very distinct from those described for HBV/HDV. Nonetheless, the mutual hepatotropism of these three viruses mediated by tissue specific factors suggests a possible overlap in usage of common hepatocyte specific host factors like NTCP.

Following establishment of the pivotal role of NTCP for HBV and HDV entry into hepatocytes, a recent study identified a role for NTCP in HCV infection (see Figure 2). Exogenous overexpression or silencing of NTCP increased or decreased HCV infection *in vitro*, respectively [114]. Unlike HBV, however, no direct interaction between HCV envelope proteins and NTCP was identified. Instead, the bile acid transporter function of NTCP was found to be important for HCV entry [114]. Bile acids are known to modulate cellular antiviral responses by inhibiting interferon (IFN) type I signaling and thereby decreasing the expression of IFN-stimulated genes (ISGs) [23, 24]. NTCP was shown to regulate HCV infection by inducing the bile acid-mediated repression of ISG expression in hepatocytes, including IFITM1, IFITM2 and IFITM3 [114]. These transmembrane proteins are known to restrict the entry of several viruses, including HCV [115]. IFITM1 blocks the interaction between HCV and its receptors [116], whereas IFITM2 and IFITM3 inhibit entry at a post-endocytosis step by blocking the release of virions into the cytoplasm [117]. NTCP facilitates HCV infection by modulating innate antiviral responses via its bile acid transport function. As bile acids have been shown to enhance HCV replication [118], it is likely that NTCP expression and activity modulates HCV infection through a multimodal mechanism of action. Interestingly, MyrB-mediated inhibition of NTCP blocks the import of bile acids, which in turn stimulates the expression of ISGs, inhibiting HCV entry and infection [114]. However, it still needs to be determined whether the inhibition of NTCP-mediated bile acid entry affects the HBV life cycle through similar mechanisms as

described for HCV. The potential of NTCP-targeting antivirals to enhance innate antiviral responses and to engage the host immune system to clear infection may be a useful property for the treatment of all hepatotropic viruses, including HBV, HCV and HDV.

Conclusions

The discovery of NTCP as the first HBV/HDV receptor was a milestone in the study of the life cycle of these viruses. This landmark discovery enabled significant progress in understanding HBV/HDV entry and virus-host interactions. Moreover, based on this discovery, novel infectious model systems based on transduced cell lines stably expressing NTCP have been developed which allow detailed study of the early steps of the viral life cycle. By allowing the study of authentic infection in cell lines, these model systems will help to understand the formation and degradation of HBV cccDNA, which is a key target to achieve the ultimate goal of HBV cure. Robust human NTCP-expressing animal model systems will enable the *in vivo* validation of virus-host interactions and antiviral therapies. Moreover, NTCP has been established as an antiviral target, and several molecules targeting NTCP are in clinical development with the goal to improve current therapies in the future. The recent discovery of NTCP as a host-dependency factor in HCV infection underscores its essential role in virus-hepatocyte interactions.

Author contributions

CFE, LH, CCC, ERV, CS, TFB wrote the manuscript.

Conflicts of interest

The authors have no conflicting interests to disclose.

286 **References**

- 287 [1] World Health Organization Global health sector strategy on viral hepatitis 2016-2021, 2016.
- 288 [2] Schweitzer A, Horn J, Mikolajczyk RT, Krause G, Ott JJ (2015) Estimations of worldwide prevalence of
289 chronic hepatitis B virus infection: A systematic review of data published between 1965 and 2013.
290 Lancet 386: 1546–1555. doi:10.1016/S0140-6736(15)61412-X.
- 291 [3] WHO Global hepatitis report, 2017, 2017.
- 292 [4] Sultanik P, Pol S (2016) Hepatitis Delta Virus: Epidemiology, Natural Course and Treatment. J Infect
293 Dis Ther 4: 2–7. doi:10.4172/2332-0877.1000271.
- 294 [5] Chung RT, Baumert TF (2014) Curing Chronic Hepatitis C — The Arc of a Medical Triumph. New
295 Engl. J. Medecine 370: 1576–1578. doi:10.1056/NEJMp1401833.
- 296 [6] Werle–Lapostolle B, Bowden S, Locarnini S, Wurstthorn K, Petersen J, Lau G, Treppe C, Marcellin P,
297 Goodman Z, Delaney WE, Xiong S, Brosgart CL, Chen S, Gibbs CS, Zoulim F (2004) Persistence of
298 cccDNA during the natural history of chronic hepatitis B and decline during adefovir dipivoxil therapy.
299 Gastroenterology 126: 1750–1758. doi:10.1053/j.gastro.2004.03.018.
- 300 [7] Papatheodoridis G V., Idilman R, Dalekos GN, Buti M, Chi H, Boemmel F Van, Calleja JL, Sympsa V,
301 Goulis J, Manolakopoulos S, Loglio A, Siakavellas S, Keskin O, Gatselis N, Hansen BE, Lehretz M,
302 Revilla J De, Savvidou S, Kourikou A, Vlachogiannakos I, Galanis K, Yurdaydin C, Berg T, Colombo
303 M, Esteban R, Janssen HLA, Lampertico P (2017) The Risk of Hepatocellular Carcinoma Decreases
304 After the First 5 Years of Entecavir or Tenofovir in Caucasians With Chronic Hepatitis B. Hepatology
305 66: 1444–1453. doi:10.1002/hep.29320.
- 306 [8] Baumert TF, Jühling F, Ono A, Hoshida Y (2017) Hepatitis C-related hepatocellular carcinoma in the era
307 of new generation antivirals. BMC Med. 15: 1–10. doi:10.1186/s12916-017-0815-7.
- 308 [9] Baumert TF, Verrier ER, Nassal M, Chung RT, Zeisel MB (2015) Host-targeting agents for treatment of
309 hepatitis B virus infection. Curr. Opin. Virol. 14: 41–46. doi:10.1016/j.coviro.2015.07.009.
- 310 [10] Mailly L, Xiao F, Lupberger J, Wilson GK, Aubert P, Duong FHT, Calabrese D, Leboeuf C, Fofana I,
311 Thumann C, Bandiera S, Lütgehetmann M, Volz T, Davis C, Harris HJ, Mee CJ, Girardi E, Chane-
312 Woon-Ming B, Ericsson M, Fletcher N, Bartenschlager R, Pessaux P, Vercauteren K, Meuleman P, Villa

313 P, Kaderali L, Pfeffer S, Heim MH, Neunlist M, Zeisel MB, Dandri M, McKeating JA, Robinet E,
314 Baumert TF (2015) Clearance of persistent hepatitis C virus infection in humanized mice using a
315 claudin-1-targeting monoclonal antibody. *Nat. Biotechnol.* 33: 549–554. doi:10.1038/nbt.3179.

316 [11] Zeisel MB, Baumert TF (2017) Clinical development of hepatitis C virus host-targeting agents. *Lancet*
317 389: 674–675. doi:10.1016/S0140-6736(17)30043-0.

318 [12] Yan H, Zhong G, Xu G, He W, Jing Z, Gao Z, Huang Y, Qi Y, Peng B, Wang H, Fu L, Song M, Chen P,
319 Gao W, Ren B, Sun Y, Cai T, Feng X, Sui J, Li W (2012) Sodium taurocholate cotransporting
320 polypeptide is a functional receptor for human hepatitis B and D virus. *Elife* 1: e000049.
321 doi:10.7554/eLife.00049.

322 [13] Ni Y, Lempp FA, Mehrle S, Nkongolo S, Kaufman C, Fälth M, Stindt J, Königer C, Nassal M, Kubitz R,
323 Sülthmann H, Urban S (2014) Hepatitis B and D viruses exploit sodium taurocholate co-transporting
324 polypeptide for species-specific entry into hepatocytes. *Gastroenterology* 146: 1070–1083.
325 doi:10.1053/j.gastro.2013.12.024.

326 [14] Volz T, Allweiss L, MBarek M Ben, Warlich M, Lohse AW, Pollok JM, Alexandrov A, Urban S,
327 Petersen J, Lütgehetmann M, Dandri M (2013) The entry inhibitor Myrcludex-B efficiently blocks
328 intrahepatic virus spreading in humanized mice previously infected with hepatitis B virus. *J. Hepatol.* 58:
329 861–867. doi:10.1016/j.jhep.2012.12.008.

330 [15] Watashi K, Sluder A, Daito T, Matsunaga S, Ryo A, Nagamori S, Iwamoto M, Nakajima S, Tsukuda S,
331 Borroto-Esoda K, Sugiyama M, Tanaka Y, Kanai Y, Kusuvara H, Mizokami M, Wakita T (2014)
332 Cyclosporin A and its analogs inhibit hepatitis B virus entry into cultured hepatocytes through targeting
333 a membrane transporter, sodium taurocholate cotransporting polypeptide (NTCP). *Hepatology* 59: 1726–
334 1737. doi:10.1002/hep.26982.

335 [16] Nkongolo S, Ni Y, Lempp FA, Kaufman C, Lindner T, Esser-Nobis K, Lohmann V, Mier W, Mehrle S,
336 Urban S (2014) Cyclosporin A inhibits hepatitis B and hepatitis D virus entry by cyclophilin-
337 independent interference with the NTCP receptor. *J. Hepatol.* 60: 723–731.
338 doi:10.1016/j.jhep.2013.11.022.

339 [17] Shimura S, Watashi K, Fukano K, Peel M, Sluder A, Kawai F, Iwamoto M, Tsukuda S, Takeuchi JS,
340 Miyake T, Sugiyama M, Ogasawara Y, Park S, Tanaka Y, Kusuvara H, Mizokami M, Sureau C, Wakita

341 T (2017) Cyclosporin derivatives inhibit hepatitis B virus entry without interfering with NTCP
342 transporter activity. *J. Hepatol.* 66: 685–692. doi:10.1016/j.jhep.2016.11.009.

343 [18] Donkers JM, Zehnder B, Westen GJP Van, Kwakkenbos MJ, Ijzerman AP, Elferink RPJO, Beuers U,
344 Urban S, van de Graaf SFJ (2017) Reduced hepatitis B and D viral entry using clinically applied drugs as
345 novel inhibitors of the bile acid transporter NTCP. *Sci. Rep.* 7: 1–13. doi:10.1038/s41598-017-15338-0.

346 [19] Kaneko M, Futamura Y, Tsukuda S, Kondoh Y, Sekine T, Hirano H, Fukano K, Ohashi H, Saso W,
347 Morishita R, Matsunaga S, Kawai F, Ryo A, Park S-Y, Suzuki R, Aizaki H, Ohtani N, Sureau C, Wakita
348 T, Osada H, Watashi K (2018) Chemical array system, a platform to identify novel hepatitis B virus
349 entry inhibitors targeting sodium taurocholate cotransporting polypeptide. *Sci. Rep.* 8: 1–13.
350 doi:10.1038/s41598-018-20987-w.

351 [20] Verrier ER, Colpitts CC, Schuster C, Zeisel MB, Baumert TF (2016) Cell culture models for the
352 investigation of Hepatitis B and D Virus infection. *Viruses* 8: 1–10. doi:10.3390/v8090261.

353 [21] Döring B, Lütteke T, Geyer J, Petzinger E (2012) The SLC10 Carrier Family: Transport Functions and
354 Molecular Structure. *Curr. Top. Membr.* 70: 105–168. doi:10.1016/B978-0-12-394316-3.00004-1.

355 [22] Esteller A (2008) Physiology of bile secretion. *World J. Gastroenterol.* 14: 5641–5649.
356 doi:10.3748/wjg.14.5641.

357 [23] Podevin P, Rosmorduc O, Conti F, Calmus Y, Meier PJ, Poupon R (1999) Bile Acids Modulate the
358 Interferon Signalling Pathway. *Hepatology* 29: 1840–1847.

359 [24] Graf D, Haselow K, Münks I, Bode JG, Häussinger D (2010) Inhibition of interferon-a-induced signaling
360 by hyperosmolarity and hydrophobic bile acids. *Biol. Chem* 391: 1175–1187. doi:10.1515/BC.2010.108.

361 [25] Slijepcevic D, van de Graaf FJ (2017) Bile Acid Uptake Transporters as Targets for Therapy. *Dig. Dis.*
362 35: 251–258. doi:10.1159/000450983.

363 [26] Geyer J, Wilke T, Petzinger E (2006) The solute carrier family SLC10: more than a family of bile acid
364 transporters regarding function and phylogenetic relationships. *Naunyn-Schmiedeberg's Arch Pharmacol*
365 372: 413–431. doi:10.1007/s00210-006-0043-8.

366 [27] Hagenbuch B, Meier PJ (1994) Molecular Cloning, Chromosomal Localization, and Functional
367 Characterization of a Human Liver Na⁺/Bile Acid Cotransporter. *J. Clin. Invest.* 93: 1326–1331.

368 [28] Mukhopadhyay S, Ananthanarayanan M, Stieger B, Meier PJ, Suchy FJ, Anwer MS (1998) Sodium
369 Taurocholate Cotransporting Polypeptide Is a Serine, Threonine Phosphoprotein and Is
370 Dephosphorylated by Cyclic Adenosine Monophosphate. *Hepatology* 92: 1629–1636.

371 [29] da Silva TC, Polli JE, Swaan PW (2013) The solute carrier family 10 (SLC10): beyond bile acid
372 transport. *Mol. Aspects Med.* 34: 252–269. doi:10.1016/j.mam.2012.07.004.The.

373 [30] Hu N-J, Iwata S, Cameron AD, Drew D (2011) Crystal structure of a bacterial homologue of the bile
374 acid sodium symporter ASBT. *Nature* 478: 408–411. doi:10.1038/nature10450.

375 [31] Dawson PA (2011) Role of the Intestinal Bile Acid Transporters in Bile Acid and Drug Disposition.
376 *Handb. Exp. Pharmacol.* 201: 169–203. doi:10.1007/978-3-642-14541-4.

377 [32] Dong Z, Ekins S, Polli JE (2013) Structure Activity Relationship for FDA Approved Drugs as Inhibitors
378 of the Human Sodium Taurocholate Co-transporting Polypeptide (NTCP). *Mol Pharm.* 10: 1008–1019.
379 doi:10.1021/mp300453k.Structure.

380 [33] Chiang JYL (2003) Bile Acid Regulation of Hepatic Physiology III. Bile acids and nuclear receptors.
381 *Am J Physiol Gastrointest Liver Physiol* 284: G349–G356.

382 [34] Goodwin B, Jones SA, Price RR, Watson MA, Mckee DD, Moore LB, Galardi C, Wilson JG, Lewis
383 MC, Roth ME, Maloney PR, Willson TM, Kliewer SA (2000) A Regulatory Cascade of the Nuclear
384 Receptors FXR , SHP-1 , and LRH-1 Represses Bile Acid Biosynthesis. *Mol. Cell* 6: 517–526.

385 [35] Lu TT, Makishima M, Repa JJ, Schoonjans K, Kerr TA, Auwerx J, Mangelsdorf DJ (2000) Molecular
386 Basis for Feedback Regulation of Bile Acid Synthesis by Nuclear Receptors. *Mol. Cell* 6: 507–515.

387 [36] Jonker JW, Stedman CAM, Liddle C, Downes M (2009) Hepatobiliary ABC transporters: physiology,
388 regulation and implications for disease. *Front. Biosci.* 14: 4904–4920. doi:10.2741/3576.

389 [37] Chiang JYL (2009) Bile acids: regulation of synthesis. *J. Lipid Res.* 50: 1955–1966.
390 doi:10.1194/jlr.R900010-JLR200.

391 [38] Gerloff T, Stieger B, Hagenbuch B, Madon J, Landmann L, Roth J, Hofmann AF, Meier PJ (1998) The
392 Sister of P-glycoprotein Represents the Canalicular Bile Salt Export Pump of Mammalian Liver. *J. Biol.*
393 *Chem.* 273: 10046–10050.

394 [39] Ananthanarayanan M, Balasubramanian N, Makishima M, Mangelsdorf DJ, Suchy FJ (2001) Human

395 Bile Salt Export Pump Promoter Is Transactivated by the Farnesoid X Receptor/Bile Acid Receptor. *J.*
396 *Biol. Chem.* 276: 28857–28865. doi:10.1074/jbc.M011610200.

397 [40] Denson LA, Sturm E, Echevarria W, Zimmerman TL, Makishima M, Mangelsdorf DJ, Karpen SJ (2001)
398 The Orphan Nuclear Receptor, shp, Mediates Bile Acid–Induced Inhibition of the Rat Bile Acid
399 Transporter, ntcp. *Gastroenterology* 121: 140–147. doi:10.1053/gast.2001.25503.

400 [41] Jung D, Hagenbuch B, Fried M, Meier PJ, Kullak-Ublick GA (2004) Role of liver-enriched transcription
401 factors and nuclear receptors in regulating the human, mouse, and rat NTCP gene. *Am J Physiol*
402 *Gastrointest Liver Physiol* 286: 752–761.

403 [42] Geier A, Martin I V, Dietrich CG, Balasubramanian N, Strauch S, Suchy FJ, Gartung C, Trautwein C,
404 Ananthanarayanan M (2008) Hepatocyte nuclear factor-4a is a central transactivator of the mouse Ntcp
405 gene. *Am J Physiol Gastrointest Liver Physiol* 295: 226–233. doi:10.1152/ajpgi.00012.2008.

406 [43] Li D, Zimmerman TL, Thevananther S, Lee H-Y, Kurie JM, Karpen SJ (2002) Interleukin-1b-mediated
407 Suppression of RXR:RAR Transactivation of the Ntcp Promoter Is JNK-dependent. *J. Biol. Chem.* 277:
408 31416–31422. doi:10.1074/jbc.M204818200.

409 [44] Siewert E, Dietrich CG, Lammert F, Heinrich PC, Matern S, Gartung C, Geier A (2004) Interleukin-6
410 regulates hepatic transporters during acute-phase response. *Biochem. Biophys. Res. Commun.* 322: 232–
411 238. doi:10.1016/j.bbrc.2004.07.102.

412 [45] Le Vee M, Lecureur V, Stieger B, Fardel O (2009) Regulation of Drug Transporter Expression in
413 Human Hepatocytes Exposed to the Proinflammatory Cytokines Tumor Necrosis Factor- α or Interleukin-
414 6. *Drug Metab. Dispos.* 37: 685–693. doi:10.1124/dmd.108.023630.pump.

415 [46] Zollner G, Fickert P, Silbert D, Fuchsbichler A, Marschall H-U, Zatloukal K, Denk H, Trauner M (2003)
416 Adaptive changes in hepatobiliary transporter expression in primary biliary cirrhosis. *J. Hepatol.* 38:
417 717–727. doi:10.1016/S0.

418 [47] Kojima H, Nies AT, König J, Hagmann W, Spring H, Uemura M, Fukui H, Keppler D (2003) Changes
419 in the expression and localization of hepatocellular transporters and radixin in primary biliary cirrhosis.
420 *J. Hepatol.* 39: 693–702. doi:10.1016/S0168-8278(03)00410-0.

421 [48] Kang J, Wang J, Cheng J, Cao Z, Chen R, Li H, Liu S, Chen X, Sui J, Lu F (2017) Down-regulation of

422 NTCP expression by cyclin D1 in hepatitis B virus-related hepatocellular carcinoma has clinical
423 significance. *Oncotarget* 8: 56041–56050.

424 [49] Anwer MS (2014) Role of protein kinase C isoforms in bile formation and cholestasis. *Hepatology* 60:
425 1090–1097. doi:10.1002/hep.27088.Role.

426 [50] Grüne S, Engelking LR, Anwer MS (1993) Role of Intracellular Calcium and Protein Kinases in the
427 Activation of Hepatic Na⁺/Taurocholate Cotransport by Cyclic AMP. *J. Biol. Chem.* 268: 17734–17741.

428 [51] Webster CRL, Blanch C, Anwer MS (2002) Role of PP2B in cAMP-induced dephosphorylation and
429 translocation of NTCP. *Am J Physiol Gastrointest Liver Physiol* 283: G44–G50.

430 [52] Anwer MS, Gillin H, Mukhopadhyay S, Balasubramaniyan N, Suchy FJ, Ananthanarayanan M (2005)
431 Dephosphorylation of Ser-226 Facilitates Plasma Membrane Retention of Ntcp. *J. Biol. Chem.* 280:
432 33687–33692. doi:10.1074/jbc.M502151200.

433 [53] Ho RH, Leake BF, Roberts RL, Lee W, Kim RB (2004) Ethnicity-dependent Polymorphism in Na⁺-
434 taurocholate Cotransporting Polypeptide (SLC10A1) Reveals a Domain Critical for Bile Acid Substrate
435 Recognition. *J. Biol. Chem.* 279: 7213–7222. doi:10.1074/jbc.M305782200.

436 [54] Pan W, Song I-S, Shin H-J, Kim M-H, Choi Y-L, Lim J, Kim W-Y, Lee S-S, Shin J-G (2011) Genetic
437 polymorphisms in Na⁺-taurocholate co-transporting polypeptide (NTCP) and ileal apical sodium-
438 dependent bile acid transporter (ASBT) and ethnic comparisons of functional variants of NTCP among
439 Asian populations. *Xenobiotica* 41: 501–510. doi:10.3109/00498254.2011.555567.

440 [55] Sureau C, Guerra B, Lanford RE (1993) Role of the Large Hepatitis B Virus Envelope Protein in
441 Infectivity of the Hepatitis Delta Virion. *J. Virol.* 67: 366–372.

442 [56] Nassal M (2015) HBV cccDNA: viral persistence reservoir and key obstacle for a cure of chronic
443 hepatitis B. *Gut* 64: 1972–1984. doi:10.1136/gutjnl-2015-309809.

444 [57] Lucifora J, Protzer U (2016) Attacking hepatitis B virus cccDNA - The holy grail to hepatitis B cure. *J.*
445 *Hepatol.* 64: S41-8. doi:10.1016/j.jhep.2016.02.009.

446 [58] Rizzetto M, Hoyer B, Canese MG, Shih JW, Purcellt RH, Gerin JL (1980) Delta agent: Association of
447 delta antigen with hepatitis B surface antigen and RNA in serum of delta-infected chimpanzees. *PNAS*
448 77: 6124–6128.

449 [59] Sureau C, Negro F (2016) The hepatitis delta virus: Replication and pathogenesis. *J. Hepatol.* 64: S102–
450 S116. doi:10.1016/j.jhep.2016.02.013.

451 [60] Marsh M, Helenius A (2006) Virus Entry: Open Sesame. *Cell* 124: 729–740.
452 doi:10.1016/j.cell.2006.02.007.

453 [61] Abou Jaoude G, Sureau C (2005) Role of the Antigenic Loop of the Hepatitis B Virus Envelope Proteins
454 in Infectivity of Hepatitis Delta Virus. *J. Virol.* 79: 10460–10466. doi:10.1128/JVI.79.16.10460.

455 [62] Abou Jaoude G, Sureau C (2007) Entry of Hepatitis Delta Virus Requires the Conserved Cysteine
456 Residues of the Hepatitis B Virus Envelope Protein Antigenic Loop and Is Blocked by Inhibitors of
457 Thiol-Disulfide Exchange. *J. Virol.* 81: 13057–13066. doi:10.1128/JVI.01495-07.

458 [63] Persing DH, Varmus HE, Ganem DON (1987) The preS1 Protein of Hepatitis B Virus Is Acylated at Its
459 Amino Terminus with Myristic Acid. *J. Virol.* 61: 1672–1677.

460 [64] Gripon P, Seyec JLE, Rumin S, Guguen-Guillouzo C (1995) Myristylation of the Hepatitis B Virus
461 Large Surface Protein Is Essential for Viral Infectivity. *Virology* 213: 292–299.

462 [65] Bruss V, Hagelstein J, Gerhardt E, Galle PR (1996) Myristylation of the Large Surface Protein Is
463 Required for Hepatitis B Virus in Vitro Infectivity. *Virology* 218: 396–399.

464 [66] Gripon P, Rumin S, Urban S, Le Seyec J, Glaise D, Cannie I, Guyomard C, Lucas J, Trepo C, Guguen-
465 Guillouzo C (2002) Infection of a human hepatoma cell line by hepatitis B virus. *Proc. Natl. Acad. Sci.*
466 U. S. A. 99: 15655–60. doi:10.1073/pnas.232137699.

467 [67] Spillmann D (2001) Heparan sulfate: Anchor for viral intruders? *Biochimie* 83: 811–817.

468 [68] Bartlett AH, Park PW (2015) Proteoglycans in host–pathogen interactions: molecular mechanisms and
469 therapeutic implications. *Expert Rev Mol Med* 12: e5. doi:10.1017/S1462399409001367.

470 [69] Schulze A, Gripon P, Urban S (2007) Hepatitis B Virus Infection Initiates With a Large Surface Protein-
471 Dependent Binding to Heparan Sulfate Proteoglycans. *Hepatology* 46: 1759–1768.
472 doi:10.1002/hep.21896.

473 [70] Sureau C, Salisse J (2013) A Conformational Heparan Sulfate Binding Site Essential to Infectivity
474 Overlaps with the Conserved Hepatitis B Virus a-Determinant. *Hepatology* 57: 985–94.
475 doi:10.1002/hep.26125.

- 476 [71] Verrier ER, Colpitts CC, Bach C, Heydmann L, Weiss A, Renaud M, Durand SC, Habersetzer F,
 477 Durantel D, Abou-Jaoudé G, López Ledesma MM, Felmlee DJ, Soumillon M, Croonenborghs T, Pochet
 478 N, Nassal M, Schuster C, Brino L, Sureau C, Zeisel MB, Baumert TF (2016) A targeted functional RNA
 479 interference screen uncovers glypican 5 as an entry factor for hepatitis B and D viruses. *Hepatology* 63:
 480 35–48. doi:10.1002/hep.28013.
- 481 [72] Leistner CM, Gruen-Bernhard S, Glebe D (2008) Role of glycosaminoglycans for binding and infection
 482 of hepatitis B virus. *Cell. Microbiol.* 10: 122–133. doi:10.1111/j.1462-5822.2007.01023.x.
- 483 [73] Neurath AR, Kent SBH, Strick N, Parker K (1986) Identification and Chemical Synthesis of a Host Cell
 484 Receptor Binding Site on Hepatitis B Virus. *Cell* 46: 429–436.
- 485 [74] Neurath BAR, Strick N, Sproul P (1992) Search for Hepatitis B Virus Cell Receptors Reveals Binding
 486 Sites for Interleukin 6 on the Virus Envelope Protein. *J. Exp. Med.* 175: 461–469.
- 487 [75] Pontisso P, Ruvoletto MG, Tiribelli C, Gerlich WH, Ruop A, Alberti A (1992) The preS1 domain of
 488 hepatitis B virus and IgA cross-react in their binding to the hepatocyte surface. *J. Gen. Virol.* 73: 2041–
 489 2045.
- 490 [76] Ryu CJ, Cho D, Gripon P, Kim HS, Guguen-Guillouzo C, Hong HJ (2000) An 80-Kilodalton Protein
 491 That Binds to the Pre-S1 Domain of Hepatitis B Virus. *J. Virol.* 74: 110–116.
- 492 [77] Falco S De, Ruvoletto MG, Verdoliva A, Ruvo M, Raucci A, Marino M, Senatore S, Cassani G, Alberti
 493 A, Pontisso P, Fassina G (2001) Cloning and Expression of a Novel Hepatitis B Virus-binding Protein
 494 from HepG2 Cells. *J. Biol. Chem.* 276: 36613–36623. doi:10.1074/jbc.M102377200.
- 495 [78] Li D, Wang XZ, Ding J, Yu J-P (2005) NACA as a Potential Cellular Target of Hepatitis B Virus PreS1
 496 Protein. *Dig. Dis. Sci.* 50: 1156–1160. doi:10.1007/s10620-005-2724-4.
- 497 [79] Zhong G, Yan H, Wang H, He W, Jing Z, Qi Y, Fu L, Gao Z, Huang Y, Xu G, Feng X, Sui J, Li W
 498 (2013) Sodium Taurocholate Cotransporting Polypeptide Mediates Woolly Monkey Hepatitis B Virus
 499 Infection of Tupaia Hepatocytes. *J. Virol.* 87: 7176–7184. doi:10.1128/JVI.03533-12.
- 500 [80] Peng L, Zhao Q, Li Q, Li M, Li C, Xu T, Jing X, Zhu X, Wang Y, Li F, Liu R, Zhong C, Pan Q, Zeng B,
 501 Liao Q, Hu B, Hu Z, Huang Y, Sham P, Liu J, Xu S, Wang J, Gao Z, Wang Y (2015) The p.Ser267Phe
 502 Variant in SLC10A1 Is Associated With Resistance to Chronic Hepatitis B. *Hepatology* 61: 1251–1260.

doi:10.1002/hep.27608.

[81] Lee HW, Park HJ, Jin B, Dezhbord M, Kim DY, Han KH, Ryu WS, Kim S, Ahn SH (2017) Effect of S267F variant of NTCP on the patients with chronic hepatitis B. *Sci. Rep.* 7: 1–7. doi:10.1038/s41598-017-17959-x.

[82] An P, Zeng Z, Winkler CA (2018) The loss-of-function S267F variant in HBV receptor NTCP reduces human risk to HBV infection and disease progression. *J Infect Dis.* Jun 14: doi:10.1093/infdis/jiy355/5037696.

[83] Hu H, Liu J, Lin Y-L, Luo W, Chu Y-J, Chang C-L, Jen C-L, Lee M-H, Lu S-N, Wang L-Y, You S-L, Yang H-I, Chen C-J (2016) The rs2296651 (S267F) variant on NTCP (SLC10A1) is inversely associated with chronic hepatitis B and progression to cirrhosis and hepatocellular carcinoma in patients with chronic hepatitis B. *Gut* 65: 1514–1521. doi:10.1136/gutjnl-2015-310686.

[84] Zollner G, Wagner M, Fickert P, Silbert D, Fuchsbichler A, Zatloukal K, Denk H, Trauner M (2005) Hepatobiliary transporter expression in human hepatocellular carcinoma. *Liver Int.* 25: 367–379. doi:10.1111/j.1478-3231.2005.01033.x.

[85] Gripon P, Diot C, Theze N, Fourel I, Loreal O, Brechot C, Guguen-Guillouzo C (1988) Hepatitis B Virus Infection of Adult Human Hepatocytes Cultured in the Presence of Dimethyl Sulfoxide. *J. Virol.* 62: 4136–4143.

[86] Lempp FA, Wiedtke E, Qu B, Roques P, Chemin I, Vondran FWR, Grand R Le, Grimm D, Urban S (2017) Sodium taurocholate cotransporting polypeptide is the limiting host factor of hepatitis B virus infection in macaque and pig hepatocytes. *Hepatology* 66: 703–716.

[87] Allweiss L, Dandri M (2016) Experimental in vitro and in vivo models for the study of human hepatitis B virus infection. *J. Hepatol.* 64: S17–S31. doi:10.1016/j.jhep.2016.02.012.

[88] He W, Ren B, Mao F, Jing Z, Li Y, Liu Y, Peng B, Yan H, Qi Y, Sun Y, Guo J-T, Sui J, Wang F, Li W (2015) Hepatitis D Virus Infection of Mice Expressing Human Sodium Taurocholate co-transporting Polypeptide. *PLoS Pathog.* 11: 1–17. doi:10.1371/journal.ppat.1004840.

[89] Lempp FA, Mutz P, Lipps C, Wirth D, Bartenschlager R, Urban S (2016) Evidence that hepatitis B virus replication in mouse cells is limited by the lack of a host cell dependency factor. *J. Hepatol.* 64: 556–

530 564. doi:10.1016/j.jhep.2015.10.030.

531 [90] Burwitz BJ, Wettengel JM, Mück-Häusl MA, Ringelhan M, Ko C, Festag MM, Hammond KB, Northrup
532 M, Bimber BN, Jacob T, Reed JS, Norris R, Park B, Moller-Tank S, Esser K, Greene JM, Wu HL,
533 Abdulhaqq S, Webb G, Sutton WF, Klug A, Swanson T, Legasse AW, Vu TQ, Asokan A, Haigwood
534 NL, Protzer U, Sacha JB (2017) Hepatocytic expression of human sodium-taurocholate cotransporting
535 polypeptide enables hepatitis B virus infection of macaques. *Nat. Commun.* 8: 1–10.
536 doi:10.1038/s41467-017-01953-y.

537 [91] Gripon P, Cannie I, Urban S (2005) Efficient Inhibition of Hepatitis B Virus Infection by Acylated
538 Peptides Derived from the Large Viral Surface Protein. *J. Virol.* 79: 1613–1622.
539 doi:10.1128/JVI.79.3.1613.

540 [92] Petersen J, Dandri M, Mier W, Lütgehetmann M, Volz T, von Weizsäcker F, Haberkorn U, Fischer L,
541 Pollok J-M, Erbes B, Seitz S, Urban S (2008) Prevention of hepatitis B virus infection in vivo by entry
542 inhibitors derived from the large envelope protein. *Nat. Biotechnol.* 26: 335–341. doi:10.1038/nbt1389.

543 [93] Blank A, Markert C, Hohmann N, Carls A, Mikus G, Lehr T, Alexandrov A, Haag M, Schwab M, Urban
544 S, Haefeli WE (2016) First-in-human application of the novel hepatitis B and hepatitis D virus entry
545 inhibitor myrcludex B. *J. Hepatol.* 65: 483–489. doi:10.1016/j.jhep.2016.04.013.

546 [94] Bogomolov P, Alexandrov A, Voronkova N, Macievich M, Kokina K, Petrachenkova M, Lehr T, Lempp
547 FA, Wedemeyer H, Haag M, Schwab M, Haefeli WE, Blank A, Urban S (2016) Treatment of chronic
548 hepatitis D with the entry inhibitor myrcludex B: First results of a phase Ib/IIa study. *J. Hepatol.* 65:
549 490–498. doi:10.1016/j.jhep.2016.04.016.

550 [95] Wedemeyer H, Bogomolov P, Blank A, Allweiss L, Dandri-Petersen M, Bremer B, Voronkova N,
551 Schöneweis K, Pathil A, Burhenne J, Haag M, Schwab M, Haefeli W-E, Wiesch JSZ, Alexandrov A,
552 Urban S (2018) Final results of a multicenter, open-label phase 2b clinical trial to assess safety and
553 efficacy of Myrcludex B in combination with Tenofovir in patients with chronic HBV/HDV co-
554 infection. *J. Hepatol.* 68: S3.

555 [96] König A, Döring B, Mohr C, Geipel A, Geyer J, Glebe D (2014) Kinetics of the bile acid transporter and
556 hepatitis B virus receptor Na⁺/taurocholate cotransporting polypeptide (NTCP) in hepatocytes. *J.*
557 *Hepatol.* 61: 867–875. doi:10.1016/j.jhep.2014.05.018.

558 [97] Blanchet M, Sureau C, Labonté P (2014) Use of FDA approved therapeutics with hNTCP metabolic
559 inhibitory properties to impair the HDV lifecycle. *Antiviral Res.* 106: 111–115.
560 doi:10.1016/j.antiviral.2014.03.017.

561 [98] Lucifora J, Esser K, Protzer U (2013) Ezetimibe blocks hepatitis B virus infection after virus uptake into
562 hepatocytes. *Antiviral Res.* 97: 195–197. doi:10.1016/j.antiviral.2012.12.008.

563 [99] Azer SA, Stacey H (1993) Differential Effects of Cyclosporin A on the transport of bile acids by human
564 hepatocytes. *Biochem. Pharmacol.* 46: 813–819.

565 [100] Mita S, Suzuki H, Akita H, Hayashi H, Onuki R, Hofmann AF, Sugiyama Y (2006) Inhibition of Bile
566 Acid Transport across Na⁺/Taurocholate Cotransporting Polypeptide (SLC10A1) and Bile Salt Export
567 Pump (ABCB 11)-Coexpressing LLC-PK1 Cells by Cholestasis-Inducing Drugs. *Drug Metab. Dispos.*
568 34: 1575–1581. doi:10.1124/dmd.105.008748.basolateral.

569 [101] Slijepcevic D, Kaufman C, Wichers CGK, Gilgioni EH, Lempp FA, Duijst S, de Waart DR, Oude
570 Elferink RPJ, Mier W, Stieger B, Beuers U, Urban S, van de Graaf SFJ (2015) Impaired Uptake of
571 Conjugated Bile Acids and Hepatitis B Virus Pres1-Binding in Na⁺-Taurocholate Cotransporting
572 Polypeptide Knockout Mice. *Hepatology* 62: 207–219. doi:10.1002/hep.27694.

573 [102] Vaz M, Paulusma CC, Huidekoper H, Ru M De, Lim C, Koster J, Ho-Mok K, Bootsma AH, Groen AK,
574 Schaap FG, Oude Elferink RPJ, Waterham HR, Wanders RJA (2015) Sodium Taurocholate
575 Cotransporting Polypeptide (SLC10A1) Deficiency: Conjugated Hypercholanemia Without a Clear
576 Clinical Phenotype. *Hepatology* 61: 260–267. doi:10.1002/hep.27240.

577 [103] Tsukuda S, Watashi K, Hojima T, Isogawa M, Iwamoto M, Omagari K, Suzuki R, Aizaki H, Kojima S,
578 Sugiyama M, Saito A, Tanaka Y, Mizokami M, Sureau C, Wakita T (2017) A New Class of Hepatitis B
579 and D Virus Entry Inhibitors, Proanthocyanidin and Its Analogs, That Directly Act on the Viral Large
580 Surface Proteins. *Hepatology* 65: 1104–1116. doi:10.1002/hep.28952.

581 [104] Khan AG, Whidby J, Miller MT, Scarborough H, Zatorski A V, Cygan A, Price AA, Yost SA,
582 Bohannon CD, Jacob J, Grakoui A, Marcotrigiano J (2014) Structure of the Core Ectodomain of the
583 Hepatitis C Virus Envelope Glycoprotein 2. *Nature* 509: 381–384. doi:10.1038/nature13117.Structure.

584 [105] Barth H, Schäfer C, Adah MI, Zhang F, Linhardt RJ, Toyoda H, Kinoshita-Toyoda A, Toida T, van
585 Kuppevelt TH, Depla E, von Weizsäcker F, Blum HE, Baumert TF (2003) Cellular Binding of Hepatitis

586 C Virus Envelope Glycoprotein E2 Requires Cell Surface Heparan Sulfate. *J. Biol. Chem.* 278: 41003–
587 41012. doi:10.1074/jbc.M302267200.

588 [106] Shi Q, Jiang J, Luo G (2013) Syndecan-1 Serves as the Major Receptor for Attachment of Hepatitis C
589 Virus to the Surfaces of Hepatocytes. *J. Virol.* 87: 6866–6875. doi:10.1128/JVI.03475-12.

590 [107] Lefevre M, Felmlee DJ, Parnot M, Baumert TF, Schuster C (2014) Syndecan 4 Is Involved in Mediating
591 HCV Entry through Interaction with Lipoviral Particle-Associated Apolipoprotein E. *PLoS One* 9: 3–10.
592 doi:10.1371/journal.pone.0095550.

593 [108] Pileri P, Uematsu Y, Campagnoli S, Galli G, Falugi F, Petracca R, Weiner AJ, Houghton M, Rosa D,
594 Grandi G, Abrignani S (1998) Binding of Hepatitis C Virus to CD81. *Science* (80-.). 282: 938–942.

595 [109] Owen DM, Huang H, Ye J, Gale Jr. M (2010) Apolipoprotein E on Hepatitis C Virion Facilitates
596 Infection through Interaction with Low Density Lipoprotein Receptor. *Virology* 394: 99–108.
597 doi:10.1016/j.virol.2009.08.037.

598 [110] Zeisel MB, Koutsoudakis G, Schnober EK, Haberstroh A, Blum HE, Cosset F-L, Wakita T, Jaeck D,
599 Doffoel M, Royer C, Soulier E, Schvoerer E, Schuster C, Stoll-Keller F, Bartenschlager R, Pietschmann
600 T, Barth H, Baumert TF (2007) Scavenger Receptor Class B Type I Is a Key Host Factor for Hepatitis C
601 Virus Infection Required for an Entry Step Closely Linked to CD81. *Hepatology* 46: 1–3.
602 doi:10.1002/hep.21994.

603 [111] Lupberger J, Zeisel MB, Xiao F, Thumann C, Fofana I, Zona L, Davis C, Mee CJ, Turek M, Gorke S,
604 Royer C, Fischer B, Zahid MN, Lavillette D, Fresquet J, Cosset F-L, Rothenberg SM, Pietschmann T,
605 Patel AH, Pessaux P, Doffoël M, Raffelsberger W, Poch O, McKeating JA, Brino L, Baumert TF (2011)
606 EGFR and EphA2 are host factors for hepatitis C virus entry and possible targets for antiviral therapy.
607 *Nat. Med.* 17: 589–595. doi:10.1038/nm.2341.

608 [112] Farquhar MJ, Hu K, Harris HJ, Davis C, Brimacombe CL, Fletcher SJ, Baumert TF, Rappoport JZ, Balfe
609 P, McKeating JA (2012) Hepatitis C Virus Induces CD81 and Claudin-1 Endocytosis. *J. Virol.* 86:
610 4305–4316. doi:10.1128/JVI.06996-11.

611 [113] Dubuisson J, Cosset F-L (2014) Virology and cell biology of the hepatitis C virus life cycle – An update.
612 *J. Hepatol.* 61: S3–S13. doi:10.1016/j.jhep.2014.06.031.

- [114] Verrier ER, Colpitts CC, Bach C, Heydmann L, Zona L, Xiao F, Thumann C, Crouch E, Gaudin R, Sureau C, Cosset F-L, McKeating JA, Pessaux P, Hoshida Y, Schuster C, Zeisel MB, Baumert TF (2016) Solute Carrier NTCP Regulates Innate Antiviral Immune Responses Targeting Hepatitis C Virus Infection of Hepatocytes. *Cell Rep.* 17: 1357–1368. doi:10.1016/j.celrep.2016.09.084.
- [115] Smith SE, Weston S, Kellam P, Marsh M (2014) IFITM proteins — cellular inhibitors of viral entry. *Curr. Opin. Virol.* 4: 71–77. doi:10.1016/j.coviro.2013.11.004.
- [116] Wilkins C, Woodward J, Lau DT-Y, Barnes A, Joyce M, McFarlane N, McKeating J, Tyrrell DL, Gale Jr. M (2013) IFITM1 is a tight junction protein that inhibits hepatitis C virus entry. *Hepatology* 57: 461–469. doi:10.1002/hep.26066.IFITM1.
- [117] Narayana SK, Helbig KJ, McCartney EM, Eyre NS, Bull RA, Eltahla A, Lloyd AR, Beard MR (2015) The Interferon-induced Transmembrane Proteins, IFITM1, IFITM2, and IFITM3 Inhibit Hepatitis C Virus Entry. *J. Biol. Chem.* 290: 25946–25959. doi:10.1074/jbc.M115.657346.
- [118] Chang K-O, George DW (2007) Bile Acids Promote the Expression of Hepatitis C Virus in Replicon-Harboring Cells. *J. Virol.* 81: 9633–9640. doi:10.1128/JVI.00795-07.

Figure legends

Fig. 1 Model of the functional role of NTCP in hepatic bile acid transport and metabolism. Transport of bile acids from portal blood into hepatocytes *via* NTCP depends on a sodium gradient and is inhibited by CsA or ezetimibe. Secretion into the bile canaliculus *via* bile salt export pump (BSEP) in an ATP-dependent manner and synthesis from cholesterol are regulated by bile acid-mediated activation of FXR. cAMP mediates dephosphorylation and membrane translocation of NTCP. NTCP: Sodium taurocholate co-transporting polypeptide; BSEP: bile salt export pump; FXR: Farnesoid X Receptor; SHP: small heterodimer partner; CYP7A1: cholesterol 7 α -hydroxylase; BA: bile acid; TJ: tight junction; CsA: cyclosporin A; cAMP: cyclic adenosine monophosphate

Fig. 2 Model of interactions between NTCP and the entry of HBV, HDV, and HCV in hepatocytes. After initial attachment to HSPG including GPC5, HBV and HDV virions bind to the receptor NTCP through the preS1-domain of the large envelope protein. NTCP inhibitors CsA and ezetimibe block viral entry like preS1-derived MyrB and CsA-derived SCY995. NTCP modulates HCV infection by interfering with innate immune responses. Bile acids interfere with the IFN signaling pathway and thereby favor HCV entry. Inhibition of NTCP-mediated bile acid import into hepatocytes promotes inhibition of HCV entry through the upregulation of ISGs including *IFITMs*. HBV: hepatitis B virus; HCV: hepatitis C virus; HDV: hepatitis D virus; HSPG: heparan sulfate proteoglycan; GPC5: glypican-5; NTCP: Sodium taurocholate co-transporting polypeptide; MyrB: myrcludex B; CsA: cyclosporin A; SCY995: synthesized CsA derivative 995; IFN: interferon; IFNAR: IFN- α/β receptor; JAK: Janus kinase; STAT: signal transducer and activator of transcription; IRF9: Interferon regulatory factor 9; ISRE: IFN-sensitive response element; ISG: IFN-stimulated gene; IFITM: IFN-induced transmembrane protein; CLDN1: Claudin 1; CD81: cluster of differentiation 81; BA: bile acid; TJ: tight junction

Figure 1



